
Asthma Action Plan

	73 0	IIIIa <i>F</i>	CCIOII	ı ıaıı						
Name	Date of Bir	th	Date /	/	407	GREEN mea	ans Go!			
Health Care Provider	Provider's I	Phone			AAK		medicine daily			
Parent/Responsible Person	Parent's Ph	ione	School		Aor	Add RESCUE m	eans Caution! nedicine			
Additional Emergency Contact	Contact Ph	one	Last 4 Digits of	SS#		RED means Get help from	a doctor now!			
Asthma Severity (see reverse si ☐ Intermittent or Persistent: ☐ Mild ☐ Moderate ☐ Set Asthma Control ☐ Well-controlled ☐ Needs better cor	/ere ☐ Cold ☐ Stroi	s Smoke (tong odors Mss/emotions	obacco, incense old/moisture Gastroesophag	e) □ Pollen [□ Pests (rode geal reflux [als	Date of Last Flu Shot: //			
Green Zone: Go!-Take	these (CONTROL	(PREVEN	ITION) N	ledicines	EVERY [Day			
You have ALL of these: Breathing is easy No cough or wheeze Can work and play Can sleep all night Peak flow in this area: (More than 80% of Personal Best) Personal best peak flow:	Inhaled cortice Inhaled cortice Inhaled cortice Leukotriene a For asthm Fast-actin	osteroid or inhaled c	e, <u>ADD:</u>	ing β-agonist , , take	after using you puff(s) MDI with nebulizer treat by mouth spacer 15 minut	h spacer tment(s) once daily a	_ times a day _ times a day t bedtime			
Yellow Zone: Caution!	Yellow Zone: Caution!–Continue CONTROL Medicines and ADD RESCUE Medicines									
You have ANY of these: • First sign of a cold • Cough or mild wheeze • Tight chest • Problems sleeping, working, or playing Peak flow in this area: to	OR Fast-acting inh Other	haled β-agonist naled β-agonist your DOCTO	nebulizer R if you have	treatment(s)	er every ho every ho more than tw doesn't work!	o times				
Red Zone: EMERGENC	Y!–Cont	tinue COI	NTROL & F	RESCUE	Medicines	and GE	T HELP!			
You have ANY of these: Can't talk, eat, or walk well Medicine is not helping Breathing hard and fast Blue lips and fingernails Tired or lethargic	OR Fast-acting inh	haled β-agonist naled β-agonist Cal	, nebulizer	r treatment <u>e</u> while giving	r <u>every 15 minu</u> very 15 minutes	, for <u>THREE</u>				
• Ribs show Peak flow in this area:	☐ Other				OR: Call 911		mbulance			
Less than (Less than 50% of Personal Best)		or go	directly to t	he Emerg	ency Depart	ment!				
REQUIRED Healthcare Provider Signa Date: REQUIRED Responsible Person Signat		Possible side efformation of the Healthcare Provement of t	ects of rescue medi ider Initials: nt is capable and a nt is <u>not</u> approved IBLE PERSON:	pproved to self- to self-medicate		cardia, tremor, a	and nervousness. above.			
Follow up with primary doctor in 1 w	eek or:	☐ I hereby authorize a trained school employee, if available, to administer medication to the student. ☐ I hereby authorize the student to possess and self-administer medication. ☐ I hereby acknowledge that the District and its schools, employees and agents shall be immune from civil liability for acts or omissions under D.C. Law 17-107 except for criminal acts, intentional wrongdoing, gross negligence, or willful misconduct.								




Asthma Action Plan

	A5U	IIIIa <i>F</i>	ACCION	riaii			
Name	Date of Birt	:h	Date /	/	424	GREEN mea	
Health Care Provider	Provider's P	hone			ASK		medicine daily
Parent/Responsible Person	Parent's Pho	one	School		J	Add RESCUE m	leans Caution! nedicine
Additional Emergency Contact	Contact Pho	one	Last 4 Digits of	F SS#	401	RED means Get help from	a doctor now!
Asthma Severity (see reverse some persistent: ☐ Mild ☐ Moderate ☐ See Asthma Control ☐ Well-controlled ☐ Needs better co	☐ Colds ☐ Stron ☐ Stress	s □ Smoke (t g odors □ M s/emotions □	Gastroesopha	e) □ Pollen □ □ Pests (roder geal reflux □	Dust □ Anim nts, cockroache Exercise	nals	Date of Last Flu Shot:
Green Zone: Go!-Take	e these C	ONTROL	(PREVEN	ITION) M	ledicines	EVERY [Day
You have ALL of these: Breathing is easy No cough or wheeze Can work and play Can sleep all night Peak flow in this area: (More than 80% of Personal Best) Personal best peak flow:	Inhaled cortico Inhaled cortico Inhaled cortico Leukotriene an For asthm	steroid or inhaled of steroid stagonist a with exercis	e, <u>ADD:</u>	ting β-agonist ,, take_	after using you puff(s) MDI wit nebulizer trea by mouth spacer 15 minus	h spacer tment(s) once daily a	_ times a day _ times a day t bedtime
Yellow Zone: Caution	!–Continu	ie CONTI	ROL Medic	ines and	ADD RESC	CUE Med	licines
You have <u>ANY</u> of these: • First sign of a cold • Cough or mild wheeze • Tight chest • Problems sleeping, working, or playing Peak flow in this area: to_ (50%-80% of Personal Best)	OR Fast-acting inha Other	aled β-agonist , -	puff(s) M nebulize R if you have or if your rescu	r treatment(s)	every ho	ours as need	
Red Zone: EMERGENO	Y!-Cont	inue COI	NTROL &	RESCUE I	Medicines	and GE	T HELP!
You have ANY of these: Can't talk, eat, or walk well Medicine is not helping Breathing hard and fast Blue lips and fingernails Tired or lethargic	Fast-acting inh OR Fast-acting inh	aled β–agonist	, puff(s) M , nebulize	r treatment <u>ev</u>	-	, for <u>THREE</u>	
• Ribs show Peak flow in this area:	☐ Other	CANNOT C	ONTACT YO	OUR DOCT	OR: Call 911	for an aı	mbulance
Less than (Less than 50% of Personal Best)			directly to t				
REQUIRED Healthcare Provider Signa Date: REQUIRED Responsible Person Signa	nture:	Possible side eff Healthcare ProvThis studeThis stude As the RESPON:I hereby astudent.	nt is capable and a nt is <u>not</u> approved SIBLE PERSON: uthorize a trained	licines (e.g., albute approved to self-a to self-medicate. school employee	erol) include tachyondminister the med	cardia, tremor, a dicine(s) named Iminister medic	and nervousness. above.
Phone:		☐ I hereby a from civil	uthorize the stude cknowledge that tl liability for acts or Il wrongdoing, gro	he District and its omissions under	s schools, employee D.C. Law 17-107 ex	es and agents sl xcept for crimin	



Asthma Action Plan

	A5U	IIIIa <i>F</i>	ACCION	riaii			
Name	Date of Birt	:h	Date /	/	424	GREEN mea	
Health Care Provider	Provider's P	hone			ASK		medicine daily
Parent/Responsible Person	Parent's Pho	one	School		J	Add RESCUE m	leans Caution! nedicine
Additional Emergency Contact	Contact Pho	one	Last 4 Digits of	F SS#	401	RED means Get help from	a doctor now!
Asthma Severity (see reverse some persistent: ☐ Mild ☐ Moderate ☐ See Asthma Control ☐ Well-controlled ☐ Needs better co	☐ Colds ☐ Stron ☐ Stress	s □ Smoke (t g odors □ M s/emotions □	Gastroesopha	e) □ Pollen □ □ Pests (roder geal reflux □	Dust □ Anim nts, cockroache Exercise	nals	Date of Last Flu Shot:
Green Zone: Go!-Take	e these C	ONTROL	(PREVEN	ITION) M	ledicines	EVERY [Day
You have ALL of these: Breathing is easy No cough or wheeze Can work and play Can sleep all night Peak flow in this area: (More than 80% of Personal Best) Personal best peak flow:	Inhaled cortico Inhaled cortico Inhaled cortico Leukotriene an For asthm	steroid or inhaled of steroid stagonist a with exercis	e, <u>ADD:</u>	ting β-agonist ,, take_	after using you puff(s) MDI wit nebulizer trea by mouth spacer 15 minus	h spacer tment(s) once daily a	_ times a day _ times a day t bedtime
Yellow Zone: Caution	!–Continu	ie CONTI	ROL Medic	ines and	ADD RESC	CUE Med	licines
You have <u>ANY</u> of these: • First sign of a cold • Cough or mild wheeze • Tight chest • Problems sleeping, working, or playing Peak flow in this area: to_ (50%-80% of Personal Best)	OR Fast-acting inha Other	aled β-agonist , -	puff(s) M nebulize R if you have or if your rescu	r treatment(s)	every ho	ours as need	
Red Zone: EMERGENO	Y!-Cont	inue COI	NTROL &	RESCUE I	Medicines	and GE	T HELP!
You have ANY of these: Can't talk, eat, or walk well Medicine is not helping Breathing hard and fast Blue lips and fingernails Tired or lethargic	Fast-acting inh OR Fast-acting inh	aled β–agonist	, puff(s) M , nebulize	r treatment <u>ev</u>	-	, for <u>THREE</u>	
• Ribs show Peak flow in this area:	☐ Other	CANNOT C	ONTACT YO	OUR DOCT	OR: Call 911	for an aı	mbulance
Less than (Less than 50% of Personal Best)			directly to t				
REQUIRED Healthcare Provider Signa Date: REQUIRED Responsible Person Signa	nture:	Possible side eff Healthcare ProvThis studeThis stude As the RESPON:I hereby astudent.	nt is capable and a nt is <u>not</u> approved SIBLE PERSON: uthorize a trained	licines (e.g., albute approved to self-a to self-medicate. school employee	erol) include tachyondminister the med	cardia, tremor, a dicine(s) named Iminister medic	and nervousness. above.
Phone:		☐ I hereby a from civil	uthorize the stude cknowledge that tl liability for acts or Il wrongdoing, gro	he District and its omissions under	s schools, employee D.C. Law 17-107 ex	es and agents sl xcept for crimin	



Stepwise Approach for Managing Asthma in Children and Adults (from 2007 NAEPP Guidelines)

			IMPAIR	MENT	RISK			
Criteria apply to all ages unless otherwise indicated	Daytime Symptoms		ttime enings ≥5 years	Interference with normal activity	Short- acting beta- agonist use	FEV ₁ % predicted (n/a in age <5)	Exacerbations requiring oral systemic corticosteroids	
	Classification of Asthma SEVERITY: TO DETERMINE INITIATION OF LONG-TERM CONTROL THERAPY Consider severity and interval since last exacerbation when assessing risk.							Step
Severe Persistent	Throughout the day	>1x/week	Often 7x/week	Extremely limited	Several x/ day	<60%	<5: ≥2 in 6 months OR ≥4 wheezing episodes in 1 year lasting >1	<5: Step 3 5-11: Step 3 Medium-dose ICS option or Step 4 12-adult: Step 4 or 5 All ages: Consider short course OCS
Moderate Persistent	Daily	3-4x/ month	>1x/week but not nightly	Some	Daily	60-80%	day AND risk factors for per- sistent asthma	<5: Step 3 5-11: Step 3 Medium-dose ICS option 12-adult: Step 3 All ages: Consider short course OCS
Mild Persistent	>2 days/ week but not daily	1-2x/ month	3-4x/ month	Minor	>2 days/ week but not daily	>80%	5-adult: ≥2/year	Step 2
Intermittent	≤2 days/week	0	≤2x/ month	None	≤2 days/ week	>80%	0-1/year	Step 1

Classification of Consider severity	Action: In children <5, consider alternate diagnosis or adjusting therapy if no benefit seen in 4-6 weeks.							
Very Poorly Controlled	Throughout the day	≥2x/week	≥4x/week	Extremely limited	Several times/day	<60%	<5: >3/year 5-adult: ≥2/year	Step up 1-2 steps. Consider short course OCS. Reevaluate in 2 weeks. For side effects, consider alternate treatment.
Not Well Controlled	>2 days/ week	≥2x/ month	1-3x/week	Some	>2 days/ week	60-80%	<5: 2-3/year 5-adult: ≥2/year	Step up at least 1 step. Reevaluate in 2-6 weeks. For side effects, consider alternate treatment.
Well Controlled	≤2 days/ week	≤1x/ month	≤2x/ month	None	≤2 days/ week	>80%	0-1/year	Maintain current treatment. Follow-up every 1-6 months. Consider step down if well controlled for at least 3 months.

Daily Doses of common inhaled corticosteroids	Low	Fluticason MDI (mcg) Medium	e High	_	Budesoni Respules (m Medium	ıg)	Be Low	clomethas MDI (mcg) Medium	one High	Fluticasone/ Salmeterol DPI	Budesonide/ Formoterol MDI
<5 years	176	>176-352	>352	0.25-0.5	>0.5-1	>1	n/a	n/a	n/a	n/a	n/a
5-11 years	88-176	>176-352	>352	0.5	1	2	80-160	>160-320	>320	100/50 mcg 1 inhalation BID	80 mcg/4.5 mcg 2 puffs BID
12 years-adult	88-264	>264-440	>440	n/a	n/a	n/a	80-240	>240-480	>480	Dose depends on patient	Dose depends on patient

SABA: Short-acting beta-agonist LABA: Long-acting beta-agonist LTRA: Leukotriene-receptor antagonist

ICS: Inhaled corticosteroids LD-ICS: Low-dose ICS MD-ICS: Medium-dose ICS HD-ICS: High-dose ICS OCS: Oral corticosteroids

CRM: Cromolyn NCM: Nedocromil THE: Theophylline MLK: Montelukast ALT: Alternative

Step 1

SABA prn

Preferred

Step 2

LD-ICS <u>Alternative</u> <5: CRM or MLK

Preferred

5-adult: CRM, LTRA, NCM, or THE Step 3

Preferred <5: MD-ICS

5-11: EITHER LD-ICS plus LABA, LTRA or THE OR MD-ICS

12-adult: LD-ICS plus LABA **OR** MD-ICS

<u>Alternative</u>

12-adult: LD-ICS plus either LTRA, THE or Zileuton

Step 4

Preferred <5: Medium-dose ICS

plus either LABA or MLK

5-adult: MD-ICS plus LABA

<u>Alternative</u> 5-11: MD-ICS plus either

LTRA or THE **12-adult:** MD-ICS *plus*

either LTRA, THE or Zileuton

Step 5

Preferred <5: HD-ICS plus either LABA or MLK

5-11: HD-ICS plus LABA

High-dose ICS plus LABA AND consider Omalizumab for patients who have allergies

<u>Alternative</u>

5-11: HD-ICS plus either LTRA or THE

Step 6

Preferred

<5: HD-ICS plus either LABA or MLK plus OCS

5-11: HD-ICS plus LABA plus OCS

12-adult:

HD-ICS plus LABA plus OCS AND consider Omalizumab for patients who have allergies

<u>Alternative</u>

5-11: HD-ICS plus either LTRA or THE plus OCS

-Step down if possible (asthma well-controlled at least 3 months)/Step up if needed (check adherence, technique, environment, co-morbidities)